



Oregon Health & Science University
Hospital and Clinics Provider's Orders

PO9031



ADULT AMBULATORY INFUSION ORDER
Tezepelumab-ekko (TEZSPIRE)
Injection

Page 1 of 2

ACCOUNT NO.
MED. REC. NO.
NAME
BIRTHDATE

Patient Identification

ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE.

Weight: _____ kg Height: _____ cm

Allergies: _____

Diagnosis Code: _____

Treatment Start Date: _____ Patient to follow up with provider on date: _____

****This plan will expire after 365 days at which time a new order will need to be placed****

GUIDELINES FOR ORDERING

1. Send **FACE SHEET and H&P or most recent chart note.**
2. Do not discontinue systemic or inhaled corticosteroids abruptly upon initiation of therapy with tezepelumab-ekko. Decrease corticosteroids gradually, if appropriate.
3. Treat patients with pre-existing helminth infections before therapy with tezepelumab-ekko. If patients become infected while receiving tezepelumab-ekko and do not respond to anti-helminth treatment, discontinue tezepelumab until the parasitic infection resolves.
4. Avoid use of live attenuated vaccines.

MEDICATIONS:

Tezepelumab-ekko (TEZSPIRE), 210 mg, subcutaneous, ONCE every 4 weeks, for _____ doses.

HYPERSENSITIVITY MEDICATIONS:

1. NURSING COMMUNICATION – If hypersensitivity or infusion reactions develop, temporarily hold the infusion and notify provider immediately. Administer emergency medications per the Treatment Algorithm for Acute Infusion Reaction (Policy HC-PAT-133-GUD, HMC C-132). Refer to algorithm for symptom monitoring and continuously assess as grade of severity may progress.
2. diphenhydrAMINE (BENADRYL) injection, 25-50 mg, intravenous, AS NEEDED x1 dose for hypersensitivity reaction
3. EPINEPHrine HCl (ADRENALIN) injection, 0.3 mg, intramuscular, AS NEEDED x1 dose for hypersensitivity reaction
4. hydrocortisone sodium succinate (SOLU-CORTEF) injection, 100 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity reaction
5. famotidine (PEPCID) injection, 20 mg, intravenous, AS NEEDED x1 dose for hypersensitivity reaction



Oregon Health & Science University
Hospital and Clinics Provider's Orders

ADULT AMBULATORY INFUSION ORDER
Tezepelumab-ekko (TEZSPIRE)
Injection

Page 2 of 2

ACCOUNT NO.
MED. REC. NO.
NAME
BIRTHDATE

Patient Identification

ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE.

By signing below, I represent the following:

I am responsible for the care of the patient (*who is identified at the top of this form*);

I hold an active, unrestricted license to practice medicine in: Oregon _____ (*check box that corresponds with state where you provide care to patient and where you are currently licensed. Specify state if not Oregon*);

My physician license Number is # _____ (MUST BE COMPLETED TO BE A VALID PRESCRIPTION); and I am acting within my scope of practice and authorized by law to order Infusion of the medication described above for the patient identified on this form.

Provider signature: _____ **Date/Time:** _____

Printed Name: _____ **Phone:** _____ **Fax:** _____

Please check the appropriate box for the patient's preferred clinic location:



OHSUHealth
Hillsboro Medical Center
FORMERLY TUALITY HEALTHCARE

Infusion Services
364 SE 8th Ave, Medical Plaza Suite 108B
Hillsboro, OR 97123
Phone number: (503) 681-4124
Fax number: (503) 681-4120

MCMC
MID-COLUMBIA MEDICAL CENTER
A Planetree Patient-Centered Hospital

Celilo Cancer Center
1800 E 19th St
The Dalles, OR 97058
Phone number: (541) 296-7585
Fax number: (541) 296-7610