

Oregon Health & Science University Hospital and Clinics Provider's Orders



ADULT AMBULATORY INFUSION ORDER

Denosumab (PROLIA) Injection

Osteoporosis

ACCOUNT NO.
MED. REC. NO.
NAME
BIRTHDATE

Page 1 of 2

Patient Identification

ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (\checkmark) TO BE ACTIVE.

Weight :kg	Height:cm
Allergies:	
Diagnosis Code:	
Γreatment Start Date:	Patient to follow up with provider on date:

GUIDELINES FOR ORDERING

- 1. Send FACE SHEET and H&P or most recent chart note.
- 2. All patients should be prescribed daily calcium and vitamin D supplementation
- 3. In patients with severe renal impairment (creatinine clearance less than 30 mL/min), high risk of hypocalcemia, disturbances of mineral metabolism (e.g. hypoparathyroidism, thyroid surgery, parathyroid surgery, malabsorption syndromes, excision of small intestines) recommend clinical monitoring of calcium, magnesium and phosphorus levels within 14 days of Prolia injection.
- 4. Risk verses benefit regarding osteonecrosis of the jaw and hip fracture must be discussed prior to treatment.
- 5. A complete metabolic panel is recommended and a calcium level must be obtained within 60 days prior to starting treatment
- 6. Pregnancy must be ruled out prior to administration. Perform pregnancy testing in all females of reproductive potential prior to administration of Prolia.
- 7. Must complete and check the following box:
 - ☐ Provider confirms that the patient has had a recent oral or dental evaluation and/or has no contraindications to therapy related to dental issues prior to initiating therapy.

LABS:

☐ Complete metabolic panel, routine, ONCE, every visit

NURSING ORDERS:

- 1. Review previous creatinine clearance, serum calcium and albumin. If no results in past 60 days order CMP.
- 2. TREATMENT PARAMETER Pharmacist to calculate corrected calcium. Hold and contact provider for corrected calcium less than 8.4 mg/dL.
- 3. Assess for new or unusual thigh, hip, groin, or jaw pain. Inform provider if positive findings or if patient is anticipating invasive dental work.
- 4. Remind patient to take at least 500 mg elemental calcium twice daily and 400 units vitamin D daily.

MEDICATIONS:

denosumab (PROLIA) injection, 60 mg, subcutaneous, every 6 months (26 weeks) x 2 doses, Administer injection into upper arm, upper thigh, or abdomen

^{**}This plan will expire after 365 days at which time a new order will need to be placed**



Oregon Health & Science University Hospital and Clinics Provider's Orders

ADULT AMBULATORY INFUSION ORDER

Denosumab (PROLIA) Injection

Osteoporosis

ACCOUNT NO.
MED. REC. NO.
NAME
BIRTHDATE

Page 2 of 2

Patient Identification

ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE.

By signing below, I represent the following: I am responsible for the care of the patient (who is in I hold an active, unrestricted license to practice medithat corresponds with state where you provide care state if not Oregon);	dicine in: 🛭 Or	egon 🗆 (check box	
My physician license Number is # PRESCRIPTION); and I am acting within my scope medication described above for the patient identified	of practice and	BE COMPLETED TO BE A VALID authorized by law to order Infusion of the	9
Provider signature:	D	ate/Time:	
Printed Name:	_ Phone:	Fax:	

Please check the appropriate box for the patient's preferred clinic location:



Infusion Services 364 SE 8th Ave, Medical Plaza Suite 108B Hillsboro, OR 97123

Phone number: (503) 681-4124 Fax number: (503) 681-4120



Celilo Cancer Center 1800 E 19th St The Dalles, OR 97058 Phone number: (541) 296-7585 Fax number: (541) 296-7610