
 <p>Oregon Health & Science University Hospital and Clinics Provider's Orders</p> <p>PO9031</p>  <p>ADULT AMBULATORY INFUSION ORDER Iron Dextran (INFED) Infusion</p> <p>Page 1 of 3</p>	<p>ACCOUNT NO. MED. REC. NO. NAME BIRTHDATE</p> <p><i>Patient Identification</i></p>
<p>ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE.</p>	

Weight: _____ kg Height: _____ cm

Allergies: _____

Diagnosis Code: _____

Treatment Start Date: _____ Patient to follow up with provider on date: _____

****This plan will expire after 365 days at which time a new order will need to be placed****

GUIDELINES FOR ORDERING

1. Send **FACE SHEET and H&P or most recent chart note.**
2. If patient is pregnant, estimated due date is: _____.
3. Provider must order and obtain a ferritin prior to patient being scheduled for iron infusion.
Labs drawn date: _____"
Copy of ferritin must be attached.
4. Many insurance providers require a ferritin result within 90 days. If ferritin is not within 90 days of signed date then patient's insurance may deny coverage for this treatment.
5. Oral iron should be discontinued prior to administration of iron dextran.
6. Premedication is not required prior to infusion of iron dextran. If premedication is needed, such as in patients with multiple drug allergies, history of asthma, or history of reaction to iron products; consider premedication with hydrocortisone. For treatment of mild infusion reactions, consider treatment with hydrocortisone. Avoid use of diphenhydramine to be used as a premedication or treatment of mild reactions.

NURSING ORDERS:

1. TREATMENT PARAMETERS – Hold treatment and notify provider if Ferritin greater than 300 ng/mL.
2. Instruct patient to set follow up appointment with provider for follow up labs.
3. Life-threatening anaphylactic reactions have occurred. Patient should be observed for anaphylactic reaction during any iron dextran administration.
4. Follow facility policies and/or protocols for vascular access maintenance with appropriate flush solution, dec clotting (alteplase), and/or dressing changes.

MEDICATIONS:

- Iron dextran (INFED) 1000 mg in sodium chloride 0.9% 500 mL, intravenous, ONCE, over 1 hour.

AS NEEDED MEDICATIONS:

1. sodium chloride 0.9%, 500 mL, intravenous, AS NEEDED x1 dose for vein discomfort. Give concurrently with iron



Oregon Health & Science University
Hospital and Clinics Provider's Orders

ADULT AMBULATORY INFUSION ORDER
Iron Dextran (INFED) Infusion

Page 2 of 3

ACCOUNT NO.
MED. REC. NO.
NAME
BIRTHDATE

Patient Identification

ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE.

HYPERSENSITIVITY MEDICATIONS:

1. NURSING COMMUNICATION – If hypersensitivity or infusion reactions develop, temporarily hold the infusion and notify provider immediately. Administer emergency medications per the Treatment Algorithm for Acute Infusion Reaction (OHSU HC-PAT-133-GUD, HMC C-132). Refer to algorithm for symptom monitoring and continuously assess as grade of severity may progress.
2. NURSING COMMUNICATION – Avoid intravenous or oral diphenhydramine, move to next option in the algorithm. Adverse effects of diphenhydramine may overlap with IV iron adverse effects such as flushing, hypotension, tachycardia.
3. EPINEPHRINE HCl (ADRENALIN) injection, 0.3 mg, intramuscular, AS NEEDED x 1 dose for hypersensitivity or infusion reaction
4. hydrocortisone sodium succinate (SOLU-CORTEF) injection, 100 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity or infusion reaction
5. famotidine (PEPCID) injection, 20 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity or infusion reaction

By signing below, I represent the following:

I am responsible for the care of the patient (*who is identified at the top of this form*);

I hold an active, unrestricted license to practice medicine in: ☐ Oregon ☐ _____ (*check box that corresponds with state where you provide care to patient and where you are currently licensed. Specify state if not Oregon*);

My physician license Number is # _____ (MUST BE COMPLETED TO BE A VALID PRESCRIPTION); and I am acting within my scope of practice and authorized by law to order Infusion of the medication described above for the patient identified on this form.

Provider signature: _____ **Date/Time:** _____

Printed Name: _____ **Phone:** _____ **Fax:** _____



**Oregon Health & Science University
Hospital and Clinics Provider's Orders**

ADULT AMBULATORY INFUSION ORDER
Iron Dextran (INFED) Infusion

Page 3 of 3

ACCOUNT NO.
MED. REC. NO.
NAME
BIRTHDATE

Patient Identification

ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE.

☒ **Please indicate the patient's preferred clinic location below**

<input type="checkbox"/> HILLSBORO MEDICAL CENTER 364 SE 8th Ave, Medical Plaza Suite 108B, Hillsboro, OR 97123	Phone (503) 681-4124 Fax (503) 681-4120
<input type="checkbox"/> ADVENTIST HEALTH – PORTLAND Infusion Services, 10123 SE Market St, Portland, OR 97216	Phone (503) 261-6631 Fax (503) 261-6756
<input type="checkbox"/> ADVENTIST HEALTH – COLUMBIA GORGE Celilo Cancer Center, 1800 E 19th St, The Dalles, OR 97058	Phone (541) 296-7585 Fax (541) 296-7610