

## Oregon Health & Science University Hospital and Clinics Provider's Orders



ADULT AMBULATORY INFUSION ORDER

Antibiotic Therapy
(Cephalosporin, Fluoroquinolone, and Others)

Page 1 of 3

ACCOUNT NO.
MED. REC. NO.
NAME
BIRTHDATE

Patient Identification

ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK ( ✓ ) TO BE ACTIVE.

Weight:	kg	Height:	cm			
Allergies:						
Diagnosis Co	de:					
Treatment Sta	art Date:	Pa	tient to follow up	with provider on da	ate:	
**This plan \	will expire afte	r 365 days at	which time a ne	ew order will need	to be placed**	
<ol> <li>Send</li> <li>If using of the admin</li> </ol>	ng this order for	and H&P or r m to request a tom of the orded ded.	er. May use amb	home health agen	cy, specify interval a ™ pump for antibiot	
☐ CMP,	with differential Routine, ONC already drawn.	E, every	(visit)(days)(w	_ (visit)(days)(week /eeks)(months) – C	ks)(months) – Circle ircle One	One
declo 2. In the	w facility policie tting (alteplase)	), and/or dress	ing changes		ce with appropriate with 5 mL D5W be	
MEDICATIO	NS:					
Cephalo	sporins:					
□ C	eFAZolin 1 gra	m in NaCl 0.99 ms over 1 day	% 100 mL IV, ON	NCE over 20-40 min NCE over 20-40 min 51.2 mL IV, ONCE o		inuous
□ C □ c in □ C	eFEPime 2 gra eFEPime 4 gra fusion via CAD	ms in NaCl 0.9 ms over 1 day D ms over 1 day	9% 50 mL IV, ON in NaCl 0.9% 10	·		
			•	ONCE over 15-30 r , ONCE over 15-30		



# Oregon Health & Science University Hospital and Clinics Provider's Orders

ADULT AMBULATORY INFUSION ORDER

Antibiotic Therapy (Cephalosporin, Fluoroquinolone, and Others)

Page 2 of 3

ACCOUNT NO. MED. REC. NO. NAME BIRTHDATE

Patient Identification

ALL ORDERS MUST BE MARKED	IN INK WITH A CHECKMARK	<b>(√</b> )	) TO BE ACTIVE

☐ CefTRIAXone 1 gram in NaCl 0.9% 50 mL IV, ONCE over 30 minutes ☐ CefTRIAXone 2 grams in NaCl 0.9% 50 mL IV, ONCE over 30 minutes
Interval: (must check one)  □ ONCE □ Daily x doses
Fluoroquinolones:
☐ Ciprofloxacin 200 mg in NaCl 0.9% 200 mL IV, ONCE over 60 minutes ☐ Ciprofloxacin 400 mg in NaCl 0.9% 200 mL IV, ONCE over 60 minutes
<ul> <li>□ Levofloxacin 250 mg in NaCl 0.9% 50 mL IV, ONCE over 60 minutes</li> <li>□ Levofloxacin 500 mg in NaCl 0.9% 100 mL IV, ONCE over 60 minutes</li> <li>□ Levofloxacin 750 mg in NaCl 0.9% 150 mL IV, ONCE over 90 minutes</li> </ul>
Interval: (must check one)  □ ONCE □ Daily x doses
Other:
<ul> <li>□ Azithromycin 250 mg in NaCl 0.9% 250 mL IV, ONCE over 60 minutes</li> <li>□ Azithromycin 500 mg in NaCl 0.9% 250 mL IV, ONCE over 60 minutes</li> </ul>
☐ Clindamycin 600 mg in NaCl 0.9% 50 mL IV, ONCE over 30 minutes ☐ Clindamycin 900 mg in NaCl 0.9% 50 mL IV, ONCE over 30 minutes
<ul> <li>□ Doxycycline 100 mg in NaCl 0.9% 250 mL IV, ONCE over 60 minutes</li> <li>□ Doxycycline 200 mg in NaCl 0.9% 250 mL IV, ONCE over 60 minutes</li> </ul>
☐ Sulfamethoxazole/Trimethoprim 5 mg/kg = mg in <b>D5W</b> IV, ONCE over 60-90 minutes
☐ Other (drug, dose, route):(Pharmacist to confirm availability)
Interval: (must check one)  □ ONCE □ Daily x doses
FOR InfuSystem™ AMBULATORY PUMP USE (hook up at infusion location):
Duration: □ days



### Oregon Health & Science University Hospital and Clinics Provider's Orders

ADULT AMBULATORY INFUSION ORDER

Antibiotic Therapy

(Cephalosporin, Fluoroquinolone, and Others)

Page 3 of 3

ACCOUNT NO.
MED. REC. NO.
NAME
BIRTHDATE

Patient Identification

ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK ( ✓ ) TO BE ACTIVE.

#### **HYPERSENSITIVITY MEDICATIONS:**

- If hypersensitivity or infusion reactions develop, temporarily hold the infusion and notify provider immediately. Administer emergency medications per the Treatment Algorithm for Acute Infusion Reaction (Policy HC-PAT-133-GUD, Tuality C-132). Refer to algorithm for symptom monitoring and continuously assess as grade of severity may progress.
- 2. diphenhydrAMINE (BENADRYL) injection, 25-50 mg, intravenous, AS NEEDED x 1 doses for hypersensitivity reaction, Max dose 50 mg
- 3. famotidine (PEPCID) injection, 20 mg, intravenous, AS NEEDED X 1 dose for hypersensitivity reaction
- 4. hydrocortisone sodium succinate (SOLU-CORTEF) injection, 100 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity reaction
- 5. EPINEPHrine HCI (ADRENALIN) injection, 0.3 mg, intramuscular, AS NEEDED x1 dose for hypersensitivity reaction

By signing below, I represent the following: am responsible for the care of the patient (who is in hold an active, unrestricted license to practice med that corresponds with state where you provide care state if not Oregon);	licine in: 🛮 Orego	on 🗆 (check box
My physician license Number is #	of practice and au	ECOMPLETED TO BE A VALID thorized by law to order Infusion of the
Provider signature:	Date	/Time:
Printed Name:	_ Phone:	Fax:

### OLC Central Intake Nurse:

Phone: 971-262-9645 (providers only) Fax: 503-346-8058

Please check the appropriate box for the patient's preferred clinic location:



Infusion Services 364 SE 8<sup>th</sup> Ave, Medical Plaza Suite 108B Hillsboro, OR 97123

Phone number: (503) 681-4124 Fax number: (503) 681-4120



Celilo Cancer Center 1800 E 19<sup>th</sup> St

The Dalles, OR 97058

Phone number: (541) 296-7585 Fax number: (541) 296-7610