Oregon Health & Science University Hospital and Clinics Provider's Orders   OHSU Health Image 1 of 2	ACCOUNT NO. MED. REC. NO. NAME BIRTHDATE Patient Identification IN INK WITH A CHECKMARK ( ✓ ) TO BE ACTIVE.					
Weight:kg Height:						
Allergies:						
Diagnosis Code:						
Treatment Start Date: Patient to follow up with provider on date:						
**This plan will expire after 365 days at which time a new order will need to be placed**						

# **GUIDELINES FOR ORDERING**

- 1. Send FACE SHEET and H&P or most recent chart note.
- 2. This product contains aluminum. Use caution with prolonged infusions in patients with renal insufficiency.
- 3. Use with caution in patients with significant cholestasis or hepatic dysfunction.
- 4. Cupric chloride is not recommended for patients with Wilson's Disease.
- 5. A serum copper level must be obtained within 30 days prior to starting treatment.

#### LABS:

□ CMP, Routine, ONCE, every \_\_\_\_\_ (visit)(days)(weeks)(months) – Circle One

## NURSING ORDERS:

- 1. Follow facility policies and/or protocols for vascular access maintenance with appropriate flush solution, declotting (alteplase), and/or dressing changes.
- 2. TREATMENT PARAMETERS: Hold copper chloride and notify provider if serum copper greater than 80 mcg/dL prior to initial treatment.
- 3. TREATMENT PARAMETERS: Hold copper chloride and notify provider if total bilirubin greater than 1.5 x ULN, or alkaline phosphatase greater than 10 x ULN prior to initiation.

## MEDICATIONS: (must check one)

cupric chloride (COPPER) 2 mg in sodium chloride 0.9% 100 mL, IV, ONCE, over 2 hours

Interval: (must check one)

Once

	Oregon Health & Science University Hospital and Clinics Provider's Orders	ACCOUNT NO.		
OHSU	ADULT AMBULATORY INFUSION ORDER	MED. REC. NO.		
Health	Cupric Chloride (COPPER)	NAME		
	<b>Infusion</b> Page 2 of 2	BIRTHDATE		
		Patient Identification		
ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK ( $\checkmark$ ) TO BE ACTIVE.				

## HYPERSENSITIVITY MEDICATIONS:

- NURSING COMMUNICATION If hypersensitivity or infusion reactions develop, temporarily hold the infusion and notify provider immediately. Administer emergency medications per the Treatment Algorithm for Acute Infusion Reaction (OHSU HC-PAT-133-GUD, HMC C-132). Refer to algorithm for symptom monitoring and continuously assess as grade of severity may progress.
- 2. diphenhydrAMINE (BENADRYL) injection, 25-50 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity or infusion reaction
- 3. EPINEPHrine HCI (ADRENALIN) injection, 0.3 mg, intramuscular, AS NEEDED x 1 dose for hypersensitivity or infusion reaction
- 4. hydrocortisone sodium succinate (SOLU-CORTEF) injection, 100 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity or infusion reaction
- 5. famotidine (PEPCID) injection, 20 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity or infusion reaction

## By signing below, I represent the following:

I am responsible for the care of the patient (*who is identified at the top of this form*); I hold an active, unrestricted license to practice medicine in: *Oregon* (*check box that corresponds with state where you provide care to patient and where you are currently licensed. Specify state if not Oregon*);

My physician license Number is # \_\_\_\_\_\_\_ (MUST BE COMPLETED TO BE A VALID PRESCRIPTION); and I am acting within my scope of practice and authorized by law to order Infusion of the medication described above for the patient identified on this form.

Provider signature:	Date/Time:	Date/Time:	
Printed Name:	Phone:	Fax:	

#### ☑ Please indicate the patient's preferred clinic location below

HILLSBORO MEDICAL CENTER	Phone	(503) 681-4124
364 SE 8th Ave, Medical Plaza Suite 108B, Hillsboro, OR 97123	Fax	(503) 681-4120
ADVENTIST HEALTH – PORTLAND Infusion Services, 10123 SE Market St, Portland, OR 97216	Phone Fax	( )
ADVENTIST HEALTH – COLUMBIA GORGE	Phone	(541) 296-7585
Celilo Cancer Center, 1800 E 19th St, The Dalles, OR 97058	Fax	(541) 296-7610